Open Access Original Article

Age and Gender-Wise Distribution of Normochromic Normocytic Anemia Subtypes in Adults: A Cross-Sectional Study

Hana Mazhar [1] Ayesha Nayyar[2], Rabia Sadaf,[3] Fakhra Noureen[4], Nadeem Ikram[5], Masooma Jaffer[6]

[1,3,4,5,6]Akhter Saeed Medical College Rawalpindi
[2]Islamic International Medical College (IIMC) Riphah University, Rawalpindi

Abstract

Objective: To determine the frequency of the two main subtypes of normochromic normocytic anemia (NNA), anemia of chronic disease (AOCD) and mixed deficiency anemia (MDA), across different adult age groups and genders, and to compare the severity of anemia between these groups.

Methodology: This descriptive cross-sectional study was conducted at the Department of Hematology, Pakistan Railway Hospital, Rawalpindi, from September 2024 to June 2025. Using non-probability convenience sampling, 174 patients aged 15–65 years with NNA on complete blood count and peripheral blood smear were enrolled. Patients were classified as AOCD or MDA on smear morphology, confirmed by serum ferritin, vitamin B12, and folate, with C-reactive protein and erythrocyte sedimentation rate as inflammatory markers.

Results: The mean age was 42.8 ± 13.2 years, and 110 (63.2%) patients were female. AOCD was found in 136 (78.2%) and MDA in 38 (21.8%) patients. Patients with MDA were younger than those with AOCD (36.4 ± 17.1 vs 44.6 ± 11.3 years; p=0.008). MDA accounted for 65.0% of NNA in the 15–25-year group, compared with 7.4–23.3% in older groups, while AOCD peaked at 46–55 years (36.8% of AOCD cases) (p<0.001). The gender distribution did not differ between subtypes (females 62.5% vs 65.8%; p=0.71). MDA was associated with lower mean hemoglobin (8.8 ± 1.3 vs 9.5 ± 1.6 g/dL; p=0.021) and more moderate-to-severe anemia (89.5% vs 55.1%; p<0.001). Severity did not differ significantly by age group (p=0.53) or gender (p=0.23).

Conclusion: AOCD was the predominant cause of NNA and was concentrated in middle-aged adults, whereas MDA predominated among adolescents and young adults and produced more severe anemia. A normocytic blood picture in a young patient should prompt evaluation of iron, vitamin B12, and folate status rather than being attributed to chronic disease.

Keywords: Anemia; Anemia of chronic disease; Mixed deficiency anemia; Normochromic normocytic anemia; Age factors; Sex factors; Pakistan.

Address for Correspondence
Dr. Hana Mazhar
Akhter Saeed Medical College Rawalpindi
hanamazhar0313@gmail.com

Introduction

Anemia is a reduction in hemoglobin concentration or red cell mass below the range expected for an individual's age and sex; the World Health Organization (WHO) defines it in non-pregnant adults as a hemoglobin concentration below 12 g/dL in women and below 13 g/dL in men.[1] In 2021, anemia affected an estimated 1.92 billion people worldwide, with women and populations in South Asia among the most affected.[2] Anemia is classified morphologically by mean corpuscular volume (MCV) into microcytic, normocytic and macrocytic types; normochromic normocytic anemia (NNA) is characterized by an MCV of 80–100 fL with normal mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration.[3,4] Because automated red cell indices are averages, examination of the peripheral blood smear (PBS) remains essential to detect mixed or dimorphic red cell populations that the indices conceal.[5,6]

NNA has two principal etiologies in adults. In anemia of chronic disease (AOCD), also termed anemia of inflammation, cytokines such as interleukin-6 induce hepcidin, which blocks intestinal iron absorption and traps iron in macrophages and hepatocytes; erythropoietin response is blunted and red cell survival shortened, producing iron-restricted erythropoiesis despite adequate iron stores.[7–9] Mixed deficiency anemia (MDA), in contrast, results from the coexistence of iron deficiency with vitamin B12 and/or folate deficiency; the opposing effects of microcytosis and macrocytosis can yield a normal MCV, while the smear shows a dimorphic picture.[5,10] MDA usually reflects inadequate dietary

Authorship Contribution: [1]Conception and design, data collection, peripheral smear examination, [2]data analysis, and manuscript drafting [3]data analysis and manuscript drafting, [4]conception and design, [5]supervision, [6]data analysis and manuscript drafting, data analysis and manuscript drafting, All authors approved the final version and agree to be accountable for all aspects of the work

Funding Source: none
Conflict of Interest: none
Received: Mar 08, 2026
Accepted: Sept 16, 2026

intake, malabsorption or gastrointestinal disease, including *Helicobacter pylori* infection, which is associated with iron deficiency.[11,12]

Distinguishing these subtypes is clinically important because treatment differs. A serum ferritin above 100 ng/mL with low transferrin saturation favours AOCD, a ferritin below 30 ng/mL indicates absolute iron deficiency, and intermediate values are difficult to interpret.[13,14] As ferritin is an acute-phase reactant, it must be interpreted together with inflammatory markers such as C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR), alongside vitamin B12 and folate assays.[15,16]

The relative contribution of these subtypes is likely to vary with age and gender. In older adults, about one-third of anemia is attributable to chronic inflammation or chronic kidney disease[17], whereas nutritional anemia predominates among adolescents and women of reproductive age in low- and middle-income countries, including Pakistan.[11,18–20] Most hospital-based studies from Pakistan have, however, examined anemia in the elderly or within single diseases such as diabetes, hypothyroidism or systemic lupus erythematosus[21–23], and data on how the subtypes of NNA are distributed across the adult age span and between genders are lacking. This study was therefore conducted to determine the frequency of AOCD and MDA among adults with NNA across different age groups and genders, and to compare the severity of anemia between these groups.

Methodology

Study design and setting: This descriptive cross-sectional study was conducted at the Department of Hematology, Pakistan Railway Hospital (PRH), from September 2024 to June 2025.

Ethical approval: The study was approved by the Institutional Review Committee (IRC) of Islamic International Medical College, Riphah International University, Rawalpindi (Ref. No. Riphah/IIMC/IRC/24/1057, dated 12 August 2024). Written informed consent was obtained from all participants (and from a guardian for participants under 18 years).

Sample size and sampling: The sample size of 174 was calculated using Cochran's formula, $n = Z^2p(1-p)/d^2$, taking an expected prevalence of the normochromic normocytic blood picture of 13% [24], a 95% confidence level (Z = 1.96) and a 5% margin of error. Patients were recruited by non-probability convenience sampling.

Inclusion criteria: Patients aged 15–65 years of either gender attending the medicine outpatient or inpatient departments whose complete blood count (CBC) showed anemia by WHO criteria[1] with normal red cell indices (MCV 80–100 fL, MCH 27–32 pg), and whose anemia was attributable to an inflammatory or systemic chronic disease (rheumatoid arthritis, systemic lupus erythematosus, diabetes mellitus, chronic kidney disease, hypothyroidism) or to nutritional deficiency or malabsorption (poor dietary intake, celiac disease, Crohn's disease, ulcerative colitis, *H. pylori* infection).

Exclusion criteria: Pregnancy; current hematinic therapy; history of blood transfusion; chronic blood loss; malignancy; tuberculosis; vasculitis; hypopituitarism; and chronic liver disease.

Data collection and laboratory methods: A detailed history and examination were recorded on a structured proforma. Venous blood was collected in EDTA, plain and citrate tubes following standard operating procedures; samples were checked for hemolysis, clotting, adequate volume and labelling. CBC was performed on a Mindray BC-5000 hematology analyzer with daily internal quality control. A peripheral smear was prepared from each sample, stained with Leishman stain and examined under oil immersion by the principal investigator for red cell morphology. Serum ferritin, vitamin B12 and folate were measured by enzyme-linked immunosorbent assay on an RT-6000 microplate reader (Rayto), CRP by a semi-quantitative latex agglutination slide test (a result >6 mg/L taken as raised), and ESR by the Westergren method.

Operational definitions: Reference ranges were: serum ferritin 10–120 ng/mL in females and 10–240 ng/mL in males[25]; serum folate 3.56–22.40 ng/mL[25]]; serum vitamin B12 206–678 pg/mL [26]; CRP <6 mg/L; and ESR up to 15 mm/h in men and 20 mm/h in women.[27] *AOCD* was defined as NNA with a normochromic normocytic smear in a patient with a documented chronic inflammatory or systemic disease, with normal or raised serum ferritin, normal vitamin B12 and folate, and raised ESR and/or CRP. *MDA* was defined as NNA with a smear showing a mixed red cell population (hypochromic microcytes with macrocytes, with or without macro-ovalocytes and hyper

segmented neutrophils), with low serum ferritin together with low vitamin B12 and/or folate. Severity of anemia was graded using the WHO hemoglobin cut-offs of 10.0 and 7.0 g/dL as mild (Hb ≥10.0 g/dL), moderate (7.0–9.9 g/dL) or severe (<7.0 g/dL), as previously applied in adult patients.[28,29] Participants were grouped by age into 15–25, 26–35, 36–45, 46–55 and 56–65 years.

Statistical analysis: Data were entered and analyzed in SPSS version 27. Continuous variables were expressed as mean ± standard deviation and compared with the independent-samples t-test (Welch's correction where variances were unequal). Categorical variables were expressed as frequencies and percentages and compared with the chi-square test. A p-value <0.05 was considered statistically significant.

Results

A total of 174 patients with NNA were included. The mean age was 42.8 ± 13.2 years (range 16–64), and 110 (63.2%) were female. AOCD was identified in 136 (78.2%) and MDA in 38 (21.8%) patients. The demographic and hematological characteristics of the two subtypes are shown in Table 1. Patients with MDA were significantly younger and had a lower mean hemoglobin than those with AOCD, whereas MCV and MCH did not differ. The proportion of females was similar in both subtypes (p=0.71).

AOCD: anemia of chronic disease; MDA: mixed deficiency anemia; MCV: mean corpuscular volume; MCH: mean corpuscular hemoglobin. Chi-square test.

The distribution of the subtypes differed significantly across age groups (p<0.001; Table 2). MDA was concentrated in the youngest group, accounting for 13 of 20 (65.0%) patients aged 15–25 years, whereas its share was 22.5% at 26–35 years and only 7.4% at 36–45 years. AOCD became more frequent with age, with the largest share of AOCD cases (50/136, 36.8%) in the 46–55-year group. Notably, all six male patients aged 15–25 years had MDA, as did 7 of 14 (50.0%) females in this age group.

Column percentages for AOCD and MDA; chi-square test (χ^2=28.2, df=4).

Overall, 65 (37.4%) patients had mild, 91 (52.3%) moderate and 18 (10.3%) severe anemia. Severity differed significantly between subtypes (p<0.001): mild anemia was more frequent in AOCD (44.9% vs 10.5%), whereas moderate (73.7% vs 46.3%) and severe (15.8% vs 8.8%) anemia were more frequent in MDA. Severity did not differ significantly between genders (p=0.23) or across age groups (p=0.53), although one-quarter of patients aged 15–25 years had severe anemia (Table 3).

Row percentages; chi-square test. Mild: Hb ≥10.0 g/dL; moderate: 7.0–9.9 g/dL; severe: <7.0 g/dL.

Discussion

This study shows that AOCD accounted for more than

Table I: Demographic and hematological characteristics of patients by subtype of normochromic normocytic anemia. (n=174)

Characteristic	AOCD (n=136)	MDA (n=38)	Total (n=174)	p-value
Age (years), mean ± SD	44.6 ± 11.3	36.4 ± 17.1	42.8 ± 13.2	0.008
Gender, n (%)				
Female	85 (62.5)	25 (65.8)	110 (63.2)	0.71
Male	51 (37.5)	13 (34.2)	64 (36.8)	
Hemoglobin (g/dL), mean ± SD	9.5 ± 1.6	8.8 ± 1.3	9.3 ± 1.5	0.021
MCV (fL), mean ± SD	86.5 ± 5.0	87.2 ± 5.3	86.6 ± 5.1	0.49
MCH (pg), mean ± SD	29.6 ± 1.4	29.6 ± 1.5	29.6 ± 1.4	0.95

Table II: Distribution of normochromic normocytic anemia subtypes across age groups (n=174)

Age group (years)	AOCD n (%)	MDA n (%)	Total n	MDA within age group (%)	p-value
15–25	7 (5.1)	13 (34.2)	20	65.0	<0.001
26–35	31 (22.8)	9 (23.7)	40	22.5	
36–45	25 (18.4)	2 (5.3)	27	7.4	
46–55	50 (36.8)	7 (18.4)	57	12.3	
56–65	23 (16.9)	7 (18.4)	30	23.3	
Total	136 (100)	38 (100)	174	21.8	

Table III: Severity of anemia according to subtype, gender and age group. (n=174)					
Variable	Mild n (%)	Moderate n (%)	Severe n (%)	Total	p-value
Subtype					
AOCD	61 (44.9)	63 (46.3)	12 (8.8)	136	<0.001
MDA	4 (10.5)	28 (73.7)	6 (15.8)	38	
Gender					
Female	37 (33.6)	63 (57.3)	10 (9.1)	110	0.23
Male	28 (43.8)	28 (43.8)	8 (12.5)	64	
Age group (years)					
15–25	5 (25.0)	10 (50.0)	5 (25.0)	20	0.53
26–35	16 (40.0)	21 (52.5)	3 (7.5)	40	
36–45	10 (37.0)	15 (55.6)	2 (7.4)	27	
46–55	24 (42.1)	29 (50.9)	4 (7.0)	57	
56–65	10 (33.3)	16 (53.3)	4 (13.3)	30	
Total	65 (37.4)	91 (52.3)	18 (10.3)	174	

three-quarters of NNA in adults, and that the two subtypes followed distinct age patterns: MDA was concentrated among adolescents and young adults, whereas AOCD rose steadily to a peak in the sixth decade. MDA was also associated with lower hemoglobin and more moderate-to-severe anemia, while the gender distribution was similar in both subtypes.

The predominance of AOCD is consistent with the view of anemia of inflammation as the commonest anemia among chronically ill and hospitalized patients.[8] Hospital-based studies from Pakistan report a similar pattern within individual chronic diseases: among diabetic inpatients in Karachi, anemia was normocytic in 59.8% and attributable to chronic kidney disease in 44%, while mixed deficiency accounted for only 6%[21]; in primary hypothyroidism, normocytic anemia was the commonest type (38.6%)[22]; and in systemic lupus erythematosus, AOCD was the leading cause of anemia (40%).[23] Our findings extend these observations to a mixed adult population and show that, even in a setting where nutritional deficiency is common, a normocytic blood picture in an adult with chronic disease most often reflects inflammation-driven anemia.

The age pattern is the principal finding of this study. AOCD is traditionally regarded as a disease of older people; in a large survey of adults aged 65 years and above, about one-third of anemia was due to chronic inflammation or renal disease.[17] By contrast, in a nationally representative sample of Indian adults aged 60 years and above, 63.5% of anemia was nutritional and anemia of chronic disease was the commonest non-nutritional type[30], suggesting that the predominance of AOCD in our cohort partly reflects its hospital-based case mix. In our cohort, which excluded patients older than 65 years, more than half of AOCD cases were nonetheless aged 46–65 years and the mean age of AOCD patients was 44.6 years, indicating that inflammation-related anemia is already common in middle age. This is likely to reflect the early onset of diabetes, chronic kidney disease and autoimmune disease in South Asia, conditions that drive hepcidin-mediated iron restriction and, in kidney disease, erythropoietin deficiency.[7,31] Conversely, MDA dominated the 15–25-year group, where it accounted for two-thirds of NNA. Adolescence and early adulthood are periods of high micronutrient demand, and in low- and middle-income countries this age group is particularly vulnerable to nutritional anemia because of poor dietary diversity, menstrual losses in women and gastrointestinal infection.[11,18] In Pakistan, pooled data show anemia in more than half of non-pregnant adolescent and adult women[32], and the National Nutrition Survey found vitamin B12 and folate deficiency each in about half of women of reproductive age[33], so combined deficiencies are to be expected in young adults. A high prevalence of anemia has also been reported among university students in Bangladesh, most commonly at 20–22 years of age.[34]

The observation that all six men aged 15–25 years had MDA suggests that dietary inadequacy and malabsorption are not confined to women. This is consistent with national data from India showing that three in ten rural men aged 15–54 years are anemic, with underweight men at higher risk.[35] Community data from rural India similarly show that the burden and pattern of anemia vary sharply with age and that gender differences emerge after puberty.[36] Taken together, these findings argue against assuming that NNA in a young adult is due to chronic disease; in this age group, iron, vitamin B12

and folate should be assessed even when indices are normal.

Women formed nearly two-thirds of both subtypes. Higher anemia rates in women are well documented in Pakistan and are linked to menstrual and reproductive losses, poverty, undernutrition and intra-household food inequity.[19,20,37] Among non-pregnant Pakistani women of reproductive age, iron deficiency anemia alone affects nearly one in five and is associated with household food insecurity and high parity.[38] However, the female excess in our AOCD group is better explained by its etiological mix: autoimmune diseases, which are strongly female-predominant[39], formed a substantial proportion of AOCD, whereas diabetes and chronic kidney disease were more common among men (data reported separately). Rheumatoid arthritis is itself frequently complicated by anemia, with mild anemia reported in 33–60% of patients.[40] In MDA, the female predominance is consistent with the nutritional vulnerability of women of reproductive age[19,20], whereas the similar severity of anemia in both genders indicates that, once NNA is established, its severity is driven more by etiology than by sex.

MDA produced more severe anemia than AOCD. AOCD is typically mild to moderate because erythropoiesis is restricted rather than abolished[8,9], whereas combined iron and vitamin B12 and/or folate deficiency impairs both hemoglobin synthesis and DNA synthesis in erythroid precursors.[11] This difference was significant using the cut-offs of 10.0 and 7.0 g/dL. These WHO thresholds were developed for pregnant women and young children[29] but have previously been applied to adults.[28] When the WHO grades specific to non-pregnant adults were applied (mild 11.0–11.9 g/dL in women and 11.0–12.9 g/dL in men; moderate 8.0–10.9 g/dL; severe <8.0 g/dL) [29], the difference was in the same direction but not statistically significant (p=0.22), largely because few patients met the adult criterion for mild anemia; with these grades, severity again did not differ by gender (p=0.25) or age group (p=0.07). However, mean hemoglobin, which does not depend on any grading, was significantly lower in MDA (8.8 vs 9.5 g/dL; p=0.021), supporting a true difference in severity. Clinically, the greater severity of MDA, together with its normal red cell indices, means that it can easily be missed if PBS examination and micronutrient assays are omitted.[5,6]

Strengths and limitations: The strengths of this study include the combination of PBS morphology with biochemical confirmation (ferritin, vitamin B12, folate, CRP and ESR) for every patient, and its focus on the working-age adult population, which is under-represented in anemia research. Its limitations include the cross-sectional design, which precludes causal inference; non-probability convenience sampling from a single hospital laboratory, which limits generalizability to the community; and the restriction of the sample to patients whose anemia was attributable to the pre-specified chronic or nutritional conditions, so the figures represent the distribution of subtypes within NNA rather than the population prevalence of NNA. The MDA group was small, particularly in some age strata, the upper age limit of 65 years excluded the elderly, and advanced markers such as transferrin saturation, soluble transferrin receptor and hepcidin were not available to identify AOCD with coexisting iron deficiency. In addition, long-term metformin therapy is associated with vitamin B12 deficiency in more than one-third of Pakistani patients with type 2 diabetes[41], so a nutritional component within the diabetic AOCD group cannot be fully excluded. Finally, severity was graded using WHO cut-offs designed for pregnancy and early childhood; results with the adult-specific WHO grades are therefore also reported.

Conclusion

AOCD was the predominant subtype of normochromic normocytic anemia in adults and was concentrated in the middle-aged, whereas MDA predominated among adolescents and young adults of both genders and was associated with more severe anemia. Age is therefore a useful guide in the evaluation of NNA: a normocytic blood picture in a young patient should prompt peripheral smear examination and assessment of iron, vitamin B12 and folate status, while NNA in middle-aged adults should prompt a search for underlying chronic disease. Larger community-based studies including older adults and advanced iron markers are recommended.

References

1. World Health Organization. Guideline on haemoglobin cutoffs to define anaemia in individuals and populations. Geneva: WHO; 2024.
2. GBD 2021 Anaemia Collaborators. Prevalence, years lived with disability, and trends in anaemia burden by severity and cause, 1990–

2021: findings from the Global Burden of Disease Study 2021. Lancet Haematol. 2023;10(9):e713–34. doi:10.1016/S2352-3026(23)00160-6
3. Yilmaz G, Shaikh H. Normochromic normocytic anemia. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan–.
4. Brill JR, Baumgardner DJ. Normocytic anemia. Am Fam Physician. 2000;62(10):2255–64.
5. Bain BJ. Diagnosis from the blood smear. N Engl J Med. 2005;353(5):498–507. doi:10.1056/NEJMra043442
6. K T N, Prasad K, Singh BMK. Analysis of red blood cells from peripheral blood smear images for anemia detection: a methodological review. Med Biol Eng Comput. 2022;60(9):2445–62. doi:10.1007/s11517-022-02614-z
7. Ganz T, Nemeth E. Iron homeostasis in host defence and inflammation. Nat Rev Immunol. 2015;15(8):500–10. doi:10.1038/nri3863
8. Weiss G, Ganz T, Goodnough LT. Anemia of inflammation. Blood. 2019;133(1):40–50. doi:10.1182/blood-2018-06-856500
9. Ganz T. Anemia of inflammation. N Engl J Med. 2019;381(12):1148–57. doi:10.1056/NEJMra1804281
10. Isik-Balci Y, Tancer-Elci H, Bor-Kucukatay M, Kilic-Erkek O, Kilic-Toprak E, Senol H, et al. Investigation of hemorheological parameters at the diagnosis and follow up of children with iron deficiency anemia and mixed anemia. Clin Hemorheol Microcirc. 2015;60(2):179–89. doi:10.3233/CH-141811
11. Chaparro CM, Suchdev PS. Anemia epidemiology, pathophysiology, and etiology in low- and middle-income countries. Ann N Y Acad Sci. 2019;1450(1):15–31. doi:10.1111/nyas.14092
12. Hudak L, Jaraisy A, Haj S, Muhsen K. An updated systematic review and meta-analysis on the association between Helicobacter pylori infection and iron deficiency anemia. Helicobacter. 2017;22(1):e12330. doi:10.1111/hel.12330
13. Weiss G, Goodnough LT. Anemia of chronic disease. N Engl J Med. 2005;352(10):1011–23. doi:10.1056/NEJMra041809
14. Camaschella C. Iron-deficiency anemia. N Engl J Med. 2015;372(19):1832–43. doi:10.1056/NEJMra1401038
15. Pfeiffer CM, Looker AC. Laboratory methodologies for indicators of iron status: strengths, limitations, and analytical challenges. Am J Clin Nutr. 2017;106(Suppl 6):1606S–14S. doi:10.3945/ajcn.117.155887
16. Cappellini MD, Comin-Colet J, de Francisco A, Dignass A, Doehner W, Lam CS, et al. Iron deficiency across chronic inflammatory conditions: international expert opinion on definition, diagnosis, and management. Am J Hematol. 2017;92(10):1068–78. doi:10.1002/ajh.24820
17. Guralnik JM, Eisenstaedt RS, Ferrucci L, Klein HG, Woodman RC. Prevalence of anemia in persons 65 years and older in the United States: evidence for a high rate of unexplained anemia. Blood. 2004;104(8):2263–8. doi:10.1182/blood-2004-05-1812
18. Owais A, Merritt C, Lee C, Bhutta ZA. Anemia among women of reproductive age: an overview of global burden, trends, determinants, and drivers of progress in low- and middle-income countries. Nutrients. 2021;13(8):2745. doi:10.3390/nu13082745
19. Harding KL, Aguayo VM, Namirembe G, Webb P. Determinants of anemia among women and children in Nepal and Pakistan: an analysis of recent national survey data. Matern Child Nutr. 2018;14(Suppl 4):e12478. doi:10.1111/mcn.12478
20. Ali SA, Abbasi Z, Shahid B, Moin G, Hambidge KM, Krebs NF, et al. Prevalence and determinants of anemia among women of reproductive age in Thatta Pakistan: findings from a cross-sectional study. PLoS One. 2020;15(9):e0239320. doi:10.1371/journal.pone.0239320
21. Shams N, Osmani MH. Newly diagnosed anemia in admitted diabetics, frequency, etiology and associated factors. J Coll Physicians Surg Pak. 2015;25(4):242–6.
22. Ali SS, Khan MN, Zafar F, Adnan STA, Yusuf A, Hussnain M, et al. Frequency and types of anemia in primary hypothyroidism patients: a prospective observational study. Cureus. 2024;16(4):e59350. doi:10.7759/cureus.59350
23. Shaikh MA, Memon I, Ghori RA. Frequency of anaemia in patients with systemic lupus erythematosus at tertiary care hospitals. J Pak Med Assoc. 2010;60(10):822–5.
24. Alqahtani SAM. Prevalence and characteristics of thyroid abnormalities and its association with anemia in ASIR region of Saudi Arabia: a cross-sectional study. Clin Pract. 2021;11(3):494–504. doi:10.3390/clinpract11030065
25. Dahman LSB, Sumaily KM, Sabi EM, Hassan MA, Thalab AMB, Sayad AS, et al. A comparative study for measuring serum ferritin levels with three different laboratory methods: enzyme-linked immunosorbent assay versus Cobas e411 and Cobas Integra 400 methods. Diagnostics (Basel). 2022;12(2):320. doi:10.3390/diagnostics12020320
26. Jassam N, Bancroft T, Luvai A, Mbagaya W, Barth JH. Vitamin B12 reference intervals on Beckman, Roche and Siemens analytical platforms. Ann Clin Biochem. 2023;60(6):417–22. doi:10.1177/00045632231202211
27. Hameed MA, Waqas S. Physiological basis and clinical utility of erythrocyte sedimentation rate. Pak J Med Sci. 2006;22(2):214–8.
28. Oikonomou IK, Fazio VW, Remzi FH, Lopez R, Lashner BA, Shen B. Risk factors for anemia in patients with ileal pouch-anal anastomosis. Dis Colon Rectum. 2007;50(1):69–74. doi:10.1007/s10350-006-0752-6
29. World Health Organization. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System (WHO/NMH/NHD/MNM/11.1). Geneva: WHO; 2011.
30. Dhankhar A, Khobragade P, Chopra G, Banerjee J, Chien S, Petrosyan S, et al. Prevalence and causes of anemia among older adults in India: findings from wave 2 of the Harmonized Diagnostic Assessment of Dementia for the Longitudinal Aging Study in India (LASI-DAD). Int J Equity Health. 2025;24(1):315. doi:10.1186/s12939-025-02671-4
31. Babitt JL, Lin HY. Mechanisms of anemia in CKD. J Am Soc Nephrol. 2012;23(10):1631–4. doi:10.1681/ASN.2011111078
32. Mahar B, Shah T, Shaikh K, Shaikh SN, Uqaili AA, Memon KN, et al. Uncovering the hidden health burden: a systematic review and meta-analysis of iron deficiency anemia among adolescents, and pregnant women in Pakistan. J Health Popul Nutr. 2024;43(1):149. doi:10.1186/s41043-024-00643-y
33. Soofi S, Khan GN, Sadiq K, Ariff S, Habib A, Kureishy S, et al. Prevalence and possible factors associated with anaemia, and vitamin B12 and folate deficiencies in women of reproductive age in Pakistan: analysis of national-level secondary survey data. BMJ Open. 2017;7(12):e018007. doi:10.1136/bmjopen-2017-018007
34. Shill KB, Karmakar P, Kibria MG, Das A, Rahman MA, Hossain MS, et al. Prevalence of iron-deficiency anaemia among university students in Noakhali region, Bangladesh. J Health Popul Nutr. 2014;32(1):103–10.
35. Singh A, Ram S, Singh S, Tripathi P. Prevalence and determinants of anaemia among men in rural India: evidence from a nationally representative survey. PLOS Glob Public Health. 2022;2(12):e0001159. doi:10.1371/journal.pgph.0001159
36. Alvarez-Uria G, Naik PK, Midde M, Yalla PS, Pakam R. Prevalence and severity of anaemia stratified by age and gender in rural India. Anemia. 2014;2014:176182. doi:10.1155/2014/176182
37. Ashraf F, Nafees Uddin MM, Mustafa MS, Mughal ZUN, Atif Aleem S. Prevalence and factors influencing anemia in women of reproductive age visiting a tertiary care hospital (Jinnah Postgraduate Medical Center) in Karachi: a cross-sectional study. Womens Health (Lond). 2024;20:17455057241227364. doi:10.1177/17455057241227364
38. Habib MA, Raynes-Greenow C, Soofi SB, Ali N, Nausheen S, Ahmed I, et al. Prevalence and determinants of iron deficiency anemia among non-pregnant women of reproductive age in Pakistan. Asia Pac J Clin Nutr. 2018;27(1):195–203. doi:10.6133/apjcn.042017.14
39. Klein SL, Flanagan KL. Sex differences in immune responses. Nat Rev Immunol. 2016;16(10):626–38. doi:10.1038/nri.2016.90

40. Wilson A, Yu HT, Goodnough LT, Nissenson AR. Prevalence and outcomes of anemia in rheumatoid arthritis: a systematic review of the literature. Am J Med. 2004;116(Suppl 7A):50S–7S. doi:10.1016/j.amjmed.2003.12.012

41. Asghar S, Tanvir H, Riaz A, Ejaz MH, Akram M, Chowdhury Evan AM, et al. Prevalence of vitamin B12 deficiency in patients with type 2 diabetes mellitus on metformin therapy: a cross-sectional study. Cureus. 2024;16(10):e72184. doi:10.7759/cureus.72184